



After Your Bronchoscopy/ Laryngoscopy

1. Do not eat or drink until your ability to swallow and your gag reflex has returned. This may take about 2 to 8 hours. You may drink and have soft foods when your gag reflex is normal.
2. When you are able to swallow and if your throat is sore, you may gargle or use lozenges. Ice cubes also ease throat discomfort.
3. Rest your voice. You may cough, clear your throat, and breathe deeply, as usual.
4. Do not smoke for 24 hours. Smoking will irritate your throat and lung tissues. It will also cause coughing and dislodge the clot.
5. If you have any of the following symptoms, contact your doctor immediately:
 - swelling
 - prolonged or increased cough
 - bleeding
 - difficulty breathing
 - chest pain while breathing
 - dizziness
 - subcutaneous emphysema (air bubbles under the skin and a crackling sensation to the touch)

Other Instructions:

Contacts (Reach the staff members below through the NIH page operator at **301-496-1211**.)

Your Doctor:



2004

This information is prepared specifically for patients participating in clinical research at the Warren Grant Magnuson Clinical Center at the National Institutes of Health and is not necessarily applicable to individuals who are patients elsewhere. If you have questions about the information presented here, talk to a member of your healthcare team.

Questions about the Clinical Center? OCCC@cc.nih.gov

Where applicable, brand names of commercial products are provided only as illustrative examples of acceptable products, and do not imply endorsement by NIH; nor does the fact that a particular brand name product is not identified imply that such product is unsatisfactory.

National Institutes of Health
Warren Grant Magnuson Clinical Center
Bethesda, Maryland 20892